

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

O P I C

OFFICE DE LA PROPRIÉTÉ
INTELLECTUELLE DU CANADA

C I P O

CANADIAN INTELLECTUAL
PROPERTY OFFICE(12) (19) (CA) **Demande-Application**(21) (A1) **2,246,414**

(86) 1997/02/14

(87) 1997/08/21


(72) HASELOFF, James Phillip, GB

(72) HODGE, Sarah, GB

(71) BTG INTERNATIONAL LIMITED, GB

(51) Int.Cl.⁶ C12N 15/82, C12N 15/67, C12N 15/62, A01H 5/00, C12N 15/31,
C12N 15/12

(30) 1996/02/14 (9603069.7) GB

(54) **AMELIORATIONS RELATIVES A L'EXPRESSION DE GENES**(54) **IMPROVEMENTS IN OR RELATING TO GENE EXPRESSION**

(57) L'invention concerne une séquence d'acide nucléique, capable de s'exprimer dans une cellule végétale, codant pour au moins une portion utile de la protéine GAL4 se fixant à l'ADN, la séquence présentant un pourcentage en bases A/T sensiblement diminué, par comparaison avec la séquence de levures de type sauvage.

(57) Disclosed is a nucleic acid sequence, expressible in a plant cell, encoding at least an effective portion of a GAL4 DNA-binding domain, the sequence having a % A/T base content substantially reduced relative to the wild-type yeast sequence.




Industrie Canada Industry Canada

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, A01H 5/00		A1	(11) International Publication Number: WO 97/30164
			(43) International Publication Date: 21 August 1997 (21.08.97)
(21) International Application Number: PCT/GB97/00406		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 14 February 1997 (14.02.97)			
(30) Priority Data: 9603069.7 14 February 1996 (14.02.96) GB			
(71) Applicant (for all designated States except US): MEDICAL RESEARCH COUNCIL [GB/GB]; 20 Park Crescent, London W1N 4AL (GB).			
(72) Inventors; and (75) Inventors/Applicants (for US only): HASELOFF, James, Phillip [AU/GB]; Glebe Cottage, Church Lane, Comberton, Cambridgeshire CB3 7ED (GB). HODGE, Sarah [GB/GB]; 59 Panton Street, Cambridge CB2 1HL (GB).			
(74) Agent: KEITH W. NASH & CO.; 90-92 Regent Street, Cambridge CB2 1DP (GB).		<p>Published</p> <p><i>With international search report.</i></p> <p><i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	

(54) Title: IMPROVEMENTS IN OR RELATING TO GENE EXPRESSION



(57) Abstract

Disclosed is a nucleic acid sequence, expressible in a plant cell, encoding at least an effective portion of a GAL4 DNA-binding domain, the sequence having a % A/T base content substantially reduced relative to the wild-type yeast sequence.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Title: Improvements in or Relating to Gene Expression

Field of the Invention

This invention relates to a nucleic acid sequence optimised for expression in a plant cell, and to vectors and plant cells comprising the nucleic acid sequence.

Background of the Invention

In yeast there is a gene called GAL4, the product of which acts as a transcriptional activator (Johnston 1987 Microbiol. Rev. 51, 458-476). The GAL4 gene is quite large (3Kb) and the encoded polypeptide comprises an N-terminal DNA-binding domain, a C terminal domain having a transcriptional activator function, and an intervening glucose-responsive element "GRE" (GAL4 being repressed in the presence of glucose).

The GAL4 protein has been quite well characterised. Amino acid residues 1-147 of the protein bind to DNA in a sequence-specific manner (Keegan *et al* 1986 Science 231, 699-704). The transcriptional activation function is associated with two short portions of the C terminal domain (Ma & Ptashne 1987 Cell 48, 847-853). The DNA sequence to which GAL4 binds has been identified as a 17mer, which must be present as a repeat for optimal GAL4 binding (Giniger *et al*, 1985 Cell 40, 767-774), and may be referred to as a GAL4-responsive "upstream activation sequence" (UAS): GAL4 binds to the UAS 5' of a gene by means of the DNA-binding domain, and the C-terminal domain causes up-regulation of transcription of the gene.

Recently, a two-element system has been developed for directing gene expression in *Drosophila melanogaster*. Brand and Perrimon (1993 Development 118: 401-415,) randomly inserted the gene encoding the GAL4 into the *Drosophila* genome using a P-

element based vector.

A large number of stable *Drosophila* lines were generated which each express GAL4 in a particular pattern, dependent on adjacent genomic DNA sequences. A chosen target gene could then be cloned under the control of GAL4 upstream activation sequences (UAS), separately transformed, and maintained silently in the absence of GAL4. Genetic crossing of this single line with any of the library of GAL4-containing lines allowed activation of the target gene in many different tissue and cell types, and the phenotypic consequences of mis-expression, including those lethal to the organism, could be conveniently studied. In addition, the library of GAL4-containing flies has become an increasingly characterised and shared resource, and which provides a common entry point for various "reverse" genetic techniques.

A similar system operable in plants would be highly desirable. However, expression of heterologous eukaryotic genes in plants has proved highly problematical in the past (see, for example, Green Fluorescent Protein) and efficient expression of GAL4 appears equally difficult, and it has been suggested that this is due to inefficient translation of GAL4 mRNA in plants (Reichel *et al*, 1995 Plant Cell Reports 14, 773-776). Ma *et al*, (1988 Nature 334, 631-633) were able to obtain transient expression of modified, functional GAL4 derivatives in tobacco-leaf protoplasts but could not demonstrate the presence of functional, full-length wild-type GAL4. Similarly, transient expression of GAL4 derivatives has been demonstrated in maize protoplasts (McCarty *et al*, 1991 Cell 66, 895-905) and when introduced by biolistic methods, in maize aleurone tissues or embryogenic calli (Goff *et al*, 1991 Genes & Dev. 5, 298-309; Goff *et al*, 1992 Genes & Dev. 6, 864-875). Hitherto however, there have been no reports of stable, efficient expression of functional GAL4 or derivatives thereof in a plant cell.

It is an aim of the present invention to provide a novel expression system operable in plants, utilising a modified portion of the GAL4 gene.

Summary of the Invention

In a first aspect the invention provides a nucleic acid sequence, expressible in a plant cell, encoding at least an effective portion of a GAL4 DNA-binding domain, the sequence having an A/T base content substantially reduced relative to the wild-type yeast sequence.

The A/T content of the wild-type yeast sequence encoding the DNA-binding domain of GAL4 is about 59%. The % A/T base content of the sequence of the invention encoding the effective portion of the GAL4 DNA-binding domain will be understood to be substantially reduced when it is less than 50%. Preferably the A/T content is less than 45%, and more preferably less than 40%. The sequence of the invention may be made, for example, by site-directed mutagenesis, or be made *de novo* by chemical synthesis.

An "effective portion" of the DNA-binding domain is a portion sufficient to retain most (i.e. over 50%) of the DNA-binding activity of the full length DNA-binding domain. Typically the "effective portion" will comprise at least two thirds of the full length sequence of the DNA-binding domain. Conveniently the nucleic acid sequence will encode substantially all of the GAL4 DNA-binding domain (i.e. amino acid residues 1-147 of the yeast polypeptide), although a substantially smaller portion (about 75 amino acid residue) is quite adequate to retain most of the DNA-binding activity (Kraulis *et al*, 1992 Nature 356, 448-450; and Marmorstein *et al*, 1992 Nature 356, 408-414). In one particular embodiment, the sequence will comprise the nucleotide sequence shown in the 5' portion (nucleotides 1 to about 460) of Figure 1, which sequence has a substantially reduced A/T content (38%), relative to the wild-type yeast sequence, yet encodes an identical amino acid sequence.

The GAL4 DNA-binding domain has no transcriptional activation function in its own right. Thus, in preferred embodiments, the sequence encoding the effective portion of the GAL4 DNA-binding domain will be operably linked to one or more other nucleic acid sequences, which sequences may be structural (i.e. encode functional polypeptides) and/or regulatory. Typically the sequence encoding the DNA-binding domain will be operably linked (e.g. fused in-frame) to a sequence encoding a peptide or polypeptide with a regulatory function, preferably a transcriptional activator. The transcriptional activator may be the activation domain of GAL4 protein, the sequence encoding which should

preferably be optimised for expression in plants (e.g. by reducing the A/T content thereof). Alternatively, the transcriptional activator could be any one of a number of such proteins known to be active in plants, which will be well-known to those skilled in the art, such that the sequence of the invention encodes a chimeric polypeptide, comprising at least an effective portion of the GAL4 DNA-binding domain and a transcriptional activation domain. A particularly suitable transcriptional activator domain is that obtainable from herpes simplex virus (HSV) VP-16, (see Greaves and O'Hare 1989 J. Virol 63, 1641-1650; VP, being also known as VMW65). Other transcriptional activation domains include certain peptides encoded by *E. coli* genomic DNA fragments (Ma & Ptashne 1987 Cell 51, 113-119) or synthetic peptides designed to form amphiphilic α -helix (Giniger & Ptashne 1987 Nature 330, 670-672). A common feature appears to be the requirement for excess charge, either positive (Gill & Ptashne 1987 Cell 51, 121-126) or, more especially for plant activation domains, excess negative charge (Estruch *et al*, 1994 Nucl. Acids Res. 22, 3983-3989). With this in mind, the person skilled in the art could readily synthesise, or find naturally-occurring sequences, which encode peptides or polypeptides with transcriptional activation activity in plants. Such activator sequences may, in any event, be modified for optimal activity in a plant cell.

In a further aspect the invention provides a nucleic acid construct, comprising the nucleic acid sequence defined above. In one particular embodiment, the nucleic acid construct could be used as an "enhancer-trap" to "fish" for plant enhancer sequences (Cf. Sundaresan *et al*, 1995 Genes & Dev. 9, 1797-1810). In such an embodiment, the construct will preferably include right and left Ti-DNA, to allow for random, stable insertion into the genome of a plant cell host. The construct will preferably comprise a naive plant promoter sequence, by which term is meant a promoter (operable in a plant cell) which requires the presence of a suitable enhancer sequence to cause substantial levels of transcription. Such a "naive" promoter may be thought of as an "enhancer-dependent" promoter, and essentially corresponds to the TATA box region of known plant promoters. "Plant" promoters includes reference to viral and bacterial promoter sequences which are active in a plant cell (e.g. nucleotides -48 to +1 of the CaMV 35S promoter). The naive promoter is conveniently substantially adjacent to one Ti border, the promoter being in competent relationship with the sequence encoding the GAL4 DNA-binding

domain fused with a transcription activation domain operable in plants such that, should the promoter become inserted in the plant host cell genome in functional relationship with an enhancer sequence, the promoter will direct the expression of the GAL4 DNA-binding domain and the transcriptional activation domain. The construct may additionally comprise a plant selectable marker (e.g. Kanamycin resistance) for convenience.

Additionally, the construct may comprise a reporter gene active in plants (such as Green Fluorescent Protein, or β -glucuronidase "GUS") in operable linkage to a GAL4-responsive upstream activation sequence (UAS), such that the reporter gene will be expressed in response to synthesis of the GAL4 DNA-binding domain/transcriptional activator fusion protein. Preferably the reporter gene is modified GFP as disclosed in WO 96/27675, or GUS (conveniently with a nuclear localisation signal). Preferably the UAS comprises one or more repeats of the 17mer to which GAL4 binds (Giniger *et al*, 1985 Cell 40, 767-774).

As an alternative method of introducing the GAL4 binding domain/transcriptional activator sequence into a plant host cell genome, the construct may comprise Ds ("dissociation") elements, which may then be moved about the genome by the action of a transiently-expressed Ac (Activator) enzyme (e.g. Cocherel *et al*, 1996 Plant Mol. Biol. 30, 539-551).

In a further aspect, the invention provides a plant or part thereof (e.g. a plant host cell or cell line) comprising the construct defined above. Typically the construct will have become integrated into a plant cell genome and be stably maintained therein. In particular the invention provides a plurality of plants or parts thereof (such as isolated plant cells, or cell lines, or tissue cultures) comprising a library, each plant or part thereof comprising a stably maintained nucleic acid sequence encoding an effective portion of the GAL4 DNA-binding domain as defined above. Conveniently the nucleic acid sequence will be incorporated into the genome of plant cells present in the library.

The library of plants or parts thereof thus provides a very useful resource. The library may be e.g. *Arabidopsis thaliana*, or of any other plant which is routinely used in

research. Each plant or part thereof in the library may have a particular pattern of expression of the integrated reporter gene (see, for example, Sundaresan *et al.*, 1995 *Genes & Dev.* 9, 1797-1810; Klimyuk *et al.*, 1995 *Molec. Gen. Genet.* 249, 357-365). Thus, introduction of a further gene, having a GAL4-responsive UAS into the cell line, will result in expression of the introduced gene in the same temporal/spatial pattern as the reporter gene, such that researchers can express a particular gene of interest in selected tissues and/or at selected times in a predictable manner.

The invention thus provides a method of expressing a gene of interest in a known (predictable) pattern in a plant or part thereof, the method comprising: introducing the gene of interest into the plant or part thereof, said gene of interest having a GAL4-responsive upstream activation sequence (UAS), characterised in that said plant or part thereof comprises a reporter gene expressed in a known pattern under the influence of a transcriptional activator comprising an effective portion of a GAL4 DNA-binding domain encoded by a sequence in accordance with the invention defined above, such that binding of the transcriptional activator to the UAS causes transcriptional activation of the gene of interest.

The temporal/spatial pattern of expression of the introduced gene or genes of interest will then mirror that already known for the reporter gene. Typically the plant or part thereof will be *Arabidopsis thaliana*, but may be any other plant which is routinely studied (e.g. maize, rice, potato etc).

Typically the gene or genes of interest will be introduced by crossing GAL4-expressing plant cell lines with a plant cell line comprising the gene(s) of interest linked to a GAL4-responsive UAS.

The nucleic acid sequence of the invention has other useful applications, in addition to use as an "enhancer trap", or to direct the expression of a gene or genes of interest in a predictable manner. The modified GAL4 DNA-binding domain/transcriptional activator may also be used to co-ordinate the expression of a plurality of genes of interest. In many circumstances it is desirable to express simultaneously a number of genes in a plant either

for research purposes, or for agricultural/industrial motives (e.g. to study metabolic pathways, or to manipulate the synthesis of desirable products such as plant dyes or lipids).

Thus, the invention provides a method for co-ordinating the expression of a plurality of genes of interest in a plant or part thereof, each of the genes of interest being functionally associated with a GAL4-responsive UAS, characterised in that the plant or part thereof comprises a sequence in accordance with the first aspect of the invention defined above and is capable of expressing a transcriptional activator comprising an effective portion of a GAL4 DNA-binding domain, such that binding of the transcriptional activator to the UAS causes simultaneous transcriptional activation of all of the genes of interest.

Conveniently the plurality of genes of interest may all be associated, in a polycistronic arrangement, with a single UAS, which facilitates their introduction into the plant or part thereof (the genes of interest typically being genes which are not naturally found in the plant or part thereof e.g. mammalian or bacterial genes, or genes from a different plant or possibly natural plant genes which have been the subject of various modifications). Alternatively one or more genes may be operably linked to a respective UAS. The transcriptional activator may be encoded by a sequence already present in the plant or part thereof, or may be introduced simultaneously with (or subsequent to) introduction of the plurality of genes of interest.

The invention will now be further described by way of illustrative examples and with reference to the accompanying drawings, of which:

Figure 1 shows a nucleotide sequence in accordance with the invention encoding a modified GAL4 DNA-binding domain, fused in frame to a sequence encoding a modified transcriptional activation domain from HSV VP16, the vertical line denotes the end of the GAL4 DNA-binding domain;

Figure 2 is a schematic representation of a nucleic acid construct comprising a sequence in accordance with the invention;

WO 97/30164

PCT/GB97/00406

8

Figure 3 shows the nucleotide sequence of part of the nucleic acid construct represented schematically in Figure 2; and

Figures 4a-4d show micrographs of roots of plants comprising a sequence in accordance with the invention.

Examples

The inventors have used an approach similar to that described by Brand & Perrimon (1993, cited above), but adapted it for use in *Arabidopsis*, using *Agrobacterium tumefaciens* - mediated transformation to produce "enhancer trap" plant cell lines with a transcription activator comprising the GAL4 DNA-binding domain, fused to the activation domain of HSV VP16 (mGAL4-VP16).

Example 1 - Construction of mGAL4-VP16 gene

Eight oligonucleotides, which correspond to the modified sequence of the GAL4 DNA binding region, were synthesised.

GAL-5' (28-mer)

GGC AAC AAT GAA GCT ACT GTC TTC TAT C (Seq. ID No. 3)

VP16-3' (31-mer)

GGC AGA TCT ACC CAC CGT ACT CGT CAA TTC C (Seq. ID No.4)

MGAL1 (109-mer)

GGC AAG CTT GGA TCC AAC AAT GAA GCT CCT GTC CTC CAT CGA GCA
GGC CTG CGA CAT CTG CCG CCT CAA GAA GCT CAA GTG CTC CAA GGA
GAA GCC GAA GTG CGC CAA G (Seq. ID No. 5)

MGAL2 (108-mer)

TCC TCT CGA GGG AAG ATC AGG AGG AAG AGC TGC TCC AGG CGC TCC

AGG CGG GAC TCC ACT TCG GTG AGG TGG GCG CGG GTC AGC GGG GAG
CGC TTG GTT TTG GGA GAG (Seq. ID No. 6)

MGAL3 (81-mer)

TTC CCT CGA GAG GAC CTC GAC ATG ATC CTG AAA ATG GAC TCC CTC
CAG GAC ATC AAA GCC CTG CTC ACC GGC CTC TTC GTC (Seq. ID No. 7)

MGAL4 (89-mer)

GGG TGA GGG GCA TGT CGG TCT CCA CGG AGG CCA GGC GGT CGG TGA
CGG CGT CTT TGT TCA CGT TGT CCT GGA CGA AGA GGC CGG TGA GC (Seq.
ID No. 8)

MGAL5 (80-mer)

GGA GAC CGA CAT GCC CCT CAC CCT GCG CCA GCA CCG CAT CAG CGC
GAC CTC CTC CTC GGA GGA GAG CAG CAA CAA GGG CC (Seq. ID No. 9)

MGAL6 (83-mer)

AGT GGA GCT CGT CCC CCA GGC TGA CGT CGG TCG GGG GGG CCG TCG
AGA CGG TCA ACT GGC GCT GGC CCT TGT TGC TGC TCT CC (Seq. ID No. 10)

The full-length oligonucleotides were purified by electrophoresis in a 5% polyacrylamide gel containing 7M urea and 90mM Tris-borate 1mM EDTA pH8.3 (TBE) buffer. The fractionated oligonucleotides were detected by brief staining with filtered 0.05% toluidine blue dye, excised and eluted overnight at 37°C in 0.5M ammonium acetate, 0.1% SDS and 0.1mM EDTA. The eluted DNAs were precipitated and washed with ethanol, and phosphorylated using T4 polynucleotide kinase. DNAs (0.5ug each) were separately resuspended in 50mM Tris-HCl pH 9.0, 10mM MgCl₂, 10mM DTT with 1mM ATP and 5 units T4 polynucleotide kinase, and incubated for 30 mins at 37°C.

The plasmid pCMVGal65 (Cousens *et al*, 1989 EMBO J. 8, 2337-2342) was used as a source of the GAL4-VP16 sequence with unmodified codon usage. Two restriction endonuclease fragments were isolated from the plasmid. A *SacI-SacI* fragment, which

contains the sequence encoding the DNA binding element of the *Saccharomyces cerevisiae* GAL4 protein, and a *SacI-KpnI* fragment, which encodes the activation domain of the herpes simplex virus VP16 protein, were purified by fractionation and elution from a low gelling temperature (LGT) 1% agarose gel.

The 5' portion of the *mGAL4-VP16* sequence was amplified by the polymerase chain reaction (PCR), using the GAL-5' and MGAL2 oligos as primers and the *SacI-SacI GAL4* DNA fragment as a template. Primers and template were incubated under standard conditions with VENT DNA polymerase (New England Biolabs) and subjected to 30 cycles of 94°C for 30 secs, 50°C for 1 min, 72°C for 1 min. The product was cut with *XhoI* restriction endonuclease and purified after electrophoresis through a 1.5% LGT agarose gel.

The MGAL4 and MGAL5 oligonucleotides were annealed by heating to 65°C and slow cooling to room temperature. The MGAL3 and MGAL6 oligonucleotides were then added and also annealed to the mixture. The oligonucleotides were then incubated with Klenow fragment of DNA polymerase 1 and T4 DNA ligase in 50mM Tris-HCl pH 7.5, 10mM MgCl₂, 10mM DTT, 1mM dNTPs and 1mM ATP for 60 mins at 37°C. An excess of MGAL3 and MGAL6 oligonucleotides were added, and the mixture was subject to PCR amplification, as above. The product was cut with the restriction endonucleases *XhoI* and *SacI*, and purified after electrophoresis through a 1.5% LGT agarose gel.

The three gel-purified restriction fragments: (1) the GAL-5'/MGAL2 PCR-amplified product, corresponding to the 5' portion of the GAL4 DNA binding domain, (2) the MGAL3-6 annealed and amplified product, corresponding to the 3' portion of the GAL4 DNA binding domain, and (3) the *SacI-KpnI* fragment from pCMVGal65, corresponding to the VP16 activation domain; were mixed and ligated with T4 DNA ligase in 50mM Tris-HCl pH 7.5, 10mM MgCl₂, 10mM DTT and 1mM ATP at 20°C overnight. The ligated product was PCR amplified (as above) using MGAL1 and VP16-3' as primers. The product, containing the entire modified *GAL4-VP16* sequence was cut with *BamHI* and ligated to PBI121 (Jefferson *et al*, EMBO J. 6:3907, 1987) which had been cut with *SacI*, blunt ended by treatment with T4 DNA polymerase, and re-cut with *BamHI*. The

WO 97/30164

PCT/GB97/00406

11

recombinant plasmid (pBIN 35S-*mGAL4-VP16*) contained the *mGAL4-VP16* gene downstream of the constitutive 35S plant promoter.

In Figure 1, the lower DNA sequence is that of the fusion gene (Seq. ID No. 1), encoding in the 5' portion the GAL4 DNA-binding domain, and encoding in the 3' portion the transcriptional activation domain from HSV VP16. The wild-type sequence is shown above for comparison. It will be seen that the wild type GAL4 DNA-binding domain coding sequence is A/T rich (A/T% content is shown on the right for each line of sequence), and it is believed that this causes inefficient expression in plants. In particular, it is believed that the A/T rich DNA comprises one or more regions which, when transcribed, are recognized as mRNA splice sites in plant cells. The altered nucleotides are shown outlined in Figure 1. The encoded polypeptide sequence is shown below (Seq. ID No. 2). The modifications to the nucleotide sequence were carefully selected to ensure that there was no resulting change in the encoded amino acid sequence. The numbers above the sequences represent the number of the nucleotide or amino acid at that position. Certain restriction endonuclease sites are also shown.

The 35S promoter and *mGAL4-VP16* gene were excised by digestion with the restriction endonucleases *HindIII* and *EcoRI*, purified by 1% LGT agarose gel electrophoresis, and subcloned into a vector containing a GAL4-responsive *mgfp5-ER* marker gene (see Example 2 below). The resulting construct is shown in Figure 2, in which R and L represent right and left Ti plasmid border sequence. The GAL4-VP16 fusion gene is downstream of the TATA-box region (nucleotides -48 to +1) of the CaMV 35S gene, were inserted adjacent to the right Ti border. The plasmid also contains the kanamycin resistance (Kan R) selectable marker, and a reporter gene (GUS or GFP) operably linked to a synthetic GAL4-UAS promoter sequence. Certain restriction endonuclease sites are also shown.

Figure 3 shows the nucleotide sequence (Seq. ID No. 15) of the plasmid across the junctions of the Ti right border, the CaMV35S TATA box, and the start of the GAL4 DNA-binding domain coding region.

The *mGAL4-VP 16* gene was directly assayed for activity in transformed *Arabidopsis* plants by *Agrobacterium*-mediated transformation (Valvekens *et al* Proc. Natl. Acad. Sci. USA 85:5536-5540, 1988).

Example 2

Construction of GAL-GFP enhancer trap vector.

Modification of the T-DNA right border.

The plasmid pBIN19 (Bevan, M. Nuc. Acids Res. 12:8711-8721, 1984) was digested with the restriction endonuclease *Sac*II, and a 2.6 Kb fragment containing the *Agrobacterium* T-DNA right border sequence was subcloned into *Sac*II cut pGEM 5Zf (Genbank accession No. X65308; commercially available from Promega). The bacteria containing the recombinant phagemid were superinfected with helper phage to produce single strand phagemid DNA.

A synthetic mutagenic oligonucleotide ("TR+": ATA TCC TGT CAA ACA CTG GAT CCG AGC TCC AAT TCA TAG TTT AAA CTG AAG GCG GG; Seq. ID No. 11) was kinased and annealed to the purified phagemid DNA, extended on the template using T7 DNA polymerase, and ligated using T4 DNA ligase. The DNA was transformed into bacteria, and recombinant plasmids (pTr+) were screened for the insertion of new sites for the restriction endonucleases *Bam*HI, *Sma*I and *Eco*RI immediately adjacent to the T-DNA right border.

Insertion of TATA box region.

The TATA box region of the 35S promoter was PCR amplified from pBI121 using oligonucleotides complementary to the -48 region of the promoter (oligonucleotide Δ 35S*Bg*II; GGC AGA TCT TCG CAA GAC CCT TCC TCT ATA TAA GG; Seq. ID No. 12) and the downstream β -glucuronidase gene (oligonucleotide GUSS*Sna*BI; CAC ACA AAC GGT GAT ACG TA; Seq. ID No. 13). The PCR product was cut with restriction endonucleases *Bg*II and *Bam*HI and ligated to *Bam*HI cut pTr+. The resulting recombinant plasmid (pTr+ Δ 35S) contained a minimal (naive) promoter positioned

adjacent to the T-DNA right border.

Insertion of 35S promoter driven *mGAL4-VP 16* gene.

Derivatives of the GAL4 with unmodified codon usage were inserted into the pTr+ Δ 35S plasmid in unsuccessful trial experiments, and one of these constructions (the insertion of a truncated GAL4 gene derived from pMA236; Ma *et al.*, 1988 Nature 334, 631-633) had resulted in the insertion of a unique *Hind*III site immediately adjacent to the *Bam*HI site in pTr+ Δ 35S. This plasmid, pTr+ Δ GAL4, was digested with restriction endonucleases *Hind*III and *Eco*RI, and ligated with the *Hind*III-*Eco*RI fragment from pBIN 35S-*mGAL4-VP 16*, to introduce the 35S-driven *mGAL4-VP 16* gene. The *Agrobacterium* sequences from this plasmid (pTr+35S-*mGAL4-VP 16*) were then recloned into a binary plant transformation vector which contained a GAL4-responsive marker gene (see below), for testing *in planta*.

Construction of a synthetic GAL4-responsive promoter.

A *Hind*III-*Xba*I restriction endonuclease fragment containing five optimised binding sites for GAL4 was excised from the plasmid pUAST (Brand & Perrimon, 1993 Development, 118, 401-415), and ligated into *Hind*III-*Xba*I cut pBI101 (Jefferson *et al.*, 1987 EMBO J. 6, 3901-3907) upstream of the GUS gene (to make pBIN Δ UAS-GUS). A sequence corresponding to the -90 region of the 35S promoter was PCR amplified using oligonucleotides DELAS1 (GAA CTC TAG AAG CTA CTC CAC GTC CAT AAG GGA CAC ATC ACA ATC CCA CTA TCC TTC GC; Seq. ID No. 14) and GUS*Sna*BI (see above). The product was *Xba*I-*Bam*HI cut and cloned into similarly cut pBIN Δ UAS-GUS. The resulting plasmid (pBIN UAS(-90-AS1)GUS) contains a synthetic GAL4 promoter upstream of the β -glucuronidase (GUS) coding sequence. The GUS gene was then replaced with that of the green fluorescent protein (GFP).

GAL4-responsive *mgfp5-ER* gene.

We have extensively mutated the green fluorescent protein gene for optimised expression in plants. The optimised gene is called *mgfp5-ER* (Siemering *et al.*, Current Biology 6:1653-1663, 1996; WO96/27675).

The *mgfp5-ER* gene has been cloned into the plant transformation vector pBI121, and we have excised a *Bam*HI-*Sac*I fragment, which contains the coding sequence, and have inserted this into *Bam*HI-*Sac*I cut pBIN UAS(-90-AS1)GUS to replace the GUS gene. This plasmid (pBIN UAS(-90-AS1)*mgfp5-ER*) was cut with *Sac*II, and ligated with the *Sac*II fragment containing the modified T-DNA right border and *mGAL4-VP16* gene from pTr+35S-*mGAL4-VP16*, to give pBIN 35S-*GAL4-VP16*+ UAS-*mgfp5-ER* (abbreviated to pBIN35S-GAL-GFP).

Testing of the pBIN35S-GAL-GFP vector.

The pBIN 35S-GAL-GFP plasmid was introduced into *Agrobacterium tumefaciens* strain LBA4044 (Jefferson *et al.*, EMBO J. 6:3901-3907, 1987) by electroporation, followed by selection on kanamycin-containing plates. The *Agrobacterium* strain was used to transform *Arabidopsis* using the technique described below. Plants were scored for the *GAL4*-dependent induction of GFP fluorescence after transformation.

Construction of GAL-GFP enhancer trap vector

The pTR+35S-*mGAL4-VP16* plasmid was treated with *Bam*HI (to excise the 35S promoter sequences) and religated using T4 DNA ligase. This plasmid (pTR+*mGAL4-VP16*) contains the modified *GAL4-VP16* positioned with a minimal promoter adjacent to the T-DNA right border. The T-DNA sequences were then excised by digestion with *Sac*II, and ligated with similarly cut pBIN UAS(-90-AS1)*mgfp5-ER*. This produced pBIN GAL-GFP, which contains the highly active *mGAL4-VP16* transcription activator, responsive to adjacent enhancer elements, and a *GAL4*-dependent GFP gene. GFP fluorescence is produced in response to *GAL4-VP16* activation.

Enhancer trap screen.

The enhancer trap vector pBIN GAL-GFP was introduced into *Agrobacterium tumefaciens* (see details below) LBA4044, and has been used to generate over 7,500 transformed *Arabidopsis* seedlings. Transgenic seedlings have been screened directly for *GAL4*-mediated GFP expression, and seeds were collected from expressing plants. We have a library of over 250 *Arabidopsis* lines which show stable inheritable patterns of *GAL4*-

mediated GFP expression in the root.

Transactivation.

To demonstrate GAL4-mediated transactivation of a foreign gene we have generated transgenic *Arabidopsis* that contain the T-DNA sequences from pBIN UAS(-90-AS1)GUS). In the absence of GAL4-VP16, these plants do not express the GUS gene. However, when genetically crossed with lines expressing the *mGAL4-VP16* gene, the GUS gene is activated in a pattern of expression which accurately reflects that of the *mGAL4-VP16* gene from the donor parent (see Example 3 below).

Example 3

A stable transformed line (J2302) of *Arabidopsis thaliana*, which formed part of the library described above, expressed modified GFP (under the influence of the *mGAL4-VP16* activator) in the cells of the extreme root tip (see Figures 4a and 4b).

Figure 4a is a low power micrograph showing GFP-mediated epifluorescence (400nm excitation) at the root tip. Figure 4b is a higher magnification confocal micrograph (488nm excitation) showing the area of GFP-mediated fluorescence in greater detail.

The line J2302 was crossed (using standard techniques) with another *Arabidopsis* line which comprised a stable, silently-maintained GUS reporter gene operably linked to a GAL4-responsive UAS. The product of the cross, as expected, proceeded to express GUS, under the influence of the GAL4-VP16 transcriptional activator, in the same pattern as the GFP reporter gene in J2302 (i.e. at the extreme root tip). Suitably stained samples, imaged by brightfield microscopy, are shown in Figures 4c and 4d, and demonstrate GUS reporter gene activity (darkest staining) at the root tip.

This illustrates that the modified GAL4 DNA-binding domain sequence can

- (i) successfully be used in the construction of enhancer trap vectors;
- (ii) express genes of interest (as exemplified by GUS) in a predictable pattern; and
- (iii) co-ordinate the simultaneous expression of a plurality of genes of interest (as

WO 97/30164

PCT/GB97/00406

16

exemplified by GFP and GUS).

Example 4

Transformation of *Arabidopsis thaliana*.

The methods employed are based on those originally disclosed by Valvekens *et al.* 1988. Proc. Natl. Acad. Sci. 85, 5536-5540).

Media

Unless otherwise stated all procedures were carried out aseptically using sterile solutions and equipment. All media are used in standard plastic disposable petri dishes or, for later stages, Magenta GA 7 pots (Magenta Corp., USA). Hormones are dissolved in dimethyl sulfoxide (DMSO) as x1000 stocks. Hormones and antibiotics are added after autoclaving and cooling of the media to 65°C.

1. Germination medium (GM) comprised: 1x Murashige and Skoog salt mixture; 1% sucrose; 100 mg/l inositol; 1.0 mg/l thiamine; 0.5 mg/l pyridoxine; 0.5 mg/l nicotinic acid; 0.5 g/l 2-(N-morpholino)ethanesulfonic acid (MES); (adjust to pH 5.7 with 1M KOH); and 0.8% Difco Bacto agar (for solid GM medium).

1a. GM K50

As GM, but supplemented with: 50 mg/l kanamycin (Sigma)

2. Callus-inducing medium (CIM) comprised: 1x Gamborg's B5 medium; 2% glucose; 0.5 g/l MES (pH 5.7, adjusted with 1N KOH); 0.5 mg/l 2,4-D (Sigma); 0.05 mg/l kinetin (Sigma); and 0.8 % agar (for solid CIM medium).

3. Shoot-inducing medium with Vancomycin (SIM V750 K50) comprised: Gamborg's B5 medium; 2% glucose; 0.5 g/l MES (pH 5.7); 0.8% agar; 5 mg/l N6-(2-isopentenyl)adenine (2ip); 0.15 mg/l indole-3-acetic acid (IAA); 750 mg/l vancomycin; and 50 mg/l kanamycin.

3a. SIM V500 K50

As SIM, but supplemented with: 500 mg/l vancomycin; and 50 mg/l kanamycin.

4. Root-inducing medium (RIM) comprised: Gamborg's B5 medium; 2% glucose; 0.5 g/l MES (pH 5.7); 0.8% agar; 12.5 mg/l indole butyric acid (IBA); and 50 mg/l cefotaxime.

Growth of plants

- (i) Place seeds into a 15 ml polypropylene centrifuge tube.
- (ii) Add ethanol for 2 min. Remove ethanol with pipette.
- (iii) Replace with 5% commercial bleach (~0.25% available chlorine) containing one drop of NP40 per 50 ml. Leave for 15 min, shaking regularly.
- (iv) Wash seeds in sterile, distilled water for at least three times.
- (v) After last wash, add the sterile seeds to 100ml of GM in a conical flask. Grow with shaking for 2-4 weeks in a culture room at 20-25°C.

Agrobacterium-mediated transformation.

1. Day One

- (i) Using a scalpel, cut-off all the green (upper) parts of the plantlets to leave only the root systems.
- (ii) Lay out the roots on plates containing solid CIM. Gently press down on each bunch of roots to ensure that they are in contact with the surface of the agar.
- (iii) Incubate for 3 d in growth room (22°C, continual light).

2. Day Four

- (i) After ensuring that there is no visible sign of contamination, collect the callus-induced roots in an empty petri dish.
- (ii) Cut the roots into 0.5 cm explants.
- (iii) Add 3-5 ml of *Agrobacterium* culture which has been grown overnight at 28°C in Luria broth, and washed by centrifugation. Swirl with blunt-nosed forceps. Leave to co-cultivate for 2min.
- (iv) Blot the roots dry on double-thick sterile filter-paper (Whatman No.1) in a petri dish.
- (v) Place the explants onto solidified CIM medium in a petri dish. and gently press them onto the surface to ensure contact with the medium.

(vi) Incubate plates in the growth room for 2 d to allow co-cultivation of *Agrobacterium*.

3. Day Six

(i) Transfer the explants to a petri dish in 20-25 ml of water.

Agitate with blunt-nosed forceps to wash off the *agrobacteria*. The bathing medium should become somewhat turbid. Transfer the root pieces to a sieve and repeat the washing. Then lift the sieve out of buffer and drain. Press down the explants down onto the mesh to remove as much buffer and *agrobacteria* as possible. Push the semi-dried explants together into a mass.

(Custom-made, autoclavable and re-usable, sieves are used to hold the explants during *Agrobacterium* infections and washings of root explants. These sieves are made from 100 ml plastic three-cornered beakers and 100 um nylon mesh. The top 3-4 cm is cut off the beaker and the mesh held in place across the bottom of this piece by pushing a ring 2 cm high cut from just above the base of the beaker into the lower portion of the top part. The two pieces are sealed together by pushing a hot metal rod through them in several places.)

(ii) Blot small bundles of root material on double-thick sterile filter-paper in a petri dish until dry and transfer to solidified SIM V750 K50 medium, taking care that root explants are in close contact with the medium.

(iii) Incubate in growth room.

Regeneration

(i) In growth room, tiny green kanamycin resistant calli appear on the yellowish root-explants after 3 weeks.

(ii) Every 2 weeks, transfer explants to fresh SIM V750 K50. Shoots (often initially vitreous) intermittently appear from the green calli over the next several weeks.

(iii) Transfer these shoots to RIM, where roots will develop over a period of 2 to 3 weeks

(iv) After the formation of roots, transfer the plantlets to GM in vented Magenta GA7 pots or directly to soil. Ensure that the humidity is kept low to ensure good seed set.

(v) Harvest seed pods when they turn yellowish-brown and are therefore mature.

Germination and screening of F1 progeny.

To test for transformation, F1 seedlings are germinated on GM K50.

- (i) Put seeds on GM K50 (surface sterilise the seeds if necessary).
- (ii) Put petri dishes in the dark at 4°C (refrigerator) for 3-5 d, to break seed dormancy. This is not necessary if seeds were stored for more than a month.
- (iii) Incubate petri dishes in growth room for 2 weeks. Sensitive seedlings form neither roots nor leaves, and have white cotyledons. Transformed seedlings are phenotypically normal.
- (iv) Transgenic callus and shoots were screened for GFP expression using an inverted fluorescence microscope (Leitz DM-IL) fitted with a filter set (Leitz-D excitation BP355-425, dichroic 455, emission LP460) suitable for the main 395nm excitation and 509nm emission peaks of GFP. The use of a 7mm threaded extension tube with a 4x objective (EF 4/0.12) gave a greater working distance, and has allowed the convenient direct observation of tissue within inverted sealed petri dishes. A 100 Watt long wavelength hand-held UV lamp (UV Products, B100AP) was also used for routine monitoring of transgenic shoots and plants. Transgenic *Arabidopsis* F1 seedlings were grown in sterile agar culture for 5 days, and were mounted in water under glass coverslips for microscopy. The specimens were examined using a BioRad MRC-600 laser-scanning confocal microscope equipped with a krypton-argon laser and filter sets suitable for the detection of fluorescein and texas red dyes (BioRad K1/K2), and a Nikon 60x PlanApo N.A. 1.2 water immersion objective.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: BTG INTERNATIONAL LIMITED

(ii) TITLE OF INVENTION: IMPROVEMENTS IN OR RELATING TO GENE
EXPRESSION

(iii) NUMBER OF SEQUENCES: 15

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: FETHERSTONHAUGH & CO.

(B) STREET: P.O. BOX 2999, STATION D

10 (C) CITY: OTTAWA

(D) STATE: ONT

(E) COUNTRY: CANADA

(F) ZIP: K1P 5Y6

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: ASCII (text)

(vi) CURRENT APPLICATION DATA:

20 (A) APPLICATION NUMBER: CA

(B) FILING DATE: 14-FEB-1997

(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: GB 9603069.7

(B) FILING DATE: 14-FEB-1996

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: FETHERSTONHAUGH & CO.

(B) REGISTRATION NUMBER:

(C) REFERENCE/DOCKET NUMBER: 23410-576

30 (ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (613)-235-4373

(B) TELEFAX: (613)-232-8440

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 701 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION:17..694

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

10	AAGCTTGGAT CCAACA ATG AAG CTC CTG TCC TCC ATC GAG CAG GCC TGC	49
	Met Lys Leu Leu Ser Ser Ile Glu Gln Ala Cys	
	1 5 10	
	GAC ATC TGC CGC CTC AAG AAG CTC AAG TGC TCC AAG GAG AAG CCG AAG	97
	Asp Ile Cys Arg Leu Lys Lys Leu Lys Cys Ser Lys Glu Lys Pro Lys	
	15 20 25	
	TGC GCC AAG TGT CTG AAG AAC AAC TGG GAG TGT CGC TAC TCT CCC AAA	145
	Cys Ala Lys Cys Leu Lys Asn Asn Trp Glu Cys Arg Tyr Ser Pro Lys	
	30 35 40	
20	ACC AAG CGC TCC CCG CTG ACC CGC GCC CAC CTC ACC GAA GTG GAG TCC	193
	Thr Lys Arg Ser Pro Leu Thr Arg Ala His Leu Thr Glu Val Glu Ser	
	45 50 55	
	CGC CTG GAG CGC CTG GAG CAG CTC TTC CTC CTG ATC TTC CCT CGA GAG	241
	Arg Leu Glu Arg Leu Glu Gln Leu Phe Leu Leu Ile Phe Pro Arg Glu	
	60 65 70 75	
30	GAC CTC GAC ATG ATC CTG AAA ATG GAC TCC CTC CAG GAC ATC AAA GCC	289
	Asp Leu Asp Met Ile Leu Lys Met Asp Ser Leu Gln Asp Ile Lys Ala	
	80 85 90	
	CTG CTC ACC GGC CTC TTC GTC CAG GAC AAC GTG AAC AAA GAC GCC GTC	337
	Leu Leu Thr Gly Leu Phe Val Gln Asp Asn Val Asn Lys Asp Ala Val	
	95 100 105	
	ACC GAC CGC CTG GCC TCC GTG GAG ACC GAC ATG CCC CTC ACC CTG CGC	385
	Thr Asp Arg Leu Ala Ser Val Glu Thr Asp Met Pro Leu Thr Leu Arg	
	110 115 120	
40	CAG CAC CGC ATC AGC GCG ACC TCC TCC TCG GAG GAG AGC AGC AAC AAG	433
	Gln His Arg Ile Ser Ala Thr Ser Ser Ser Glu Glu Ser Ser Asn Lys	
	125 130 135	
	GGC CAG CGC CAG TTG ACC GTC TCG ACG GCC CCC CCG ACC GAC GTC AGC	481
	Gly Gln Arg Gln Leu Thr Val Ser Thr Ala Pro Pro Thr Asp Val Ser	
	140 145 150 155	
50	CTG GGG GAC GAG CTC CAC TTA GAC GGC GAG GAC GTG GCG ATG GCG CAT	529
	Leu Gly Asp Glu Leu His Leu Asp Gly Glu Asp Val Ala Met Ala His	
	160 165 170	
	GCC GAC GCG CTA GAC GAT TTC GAT CTG GAC ATG TTG GGG GAC GGG GAT	577
	Ala Asp Ala Leu Asp Asp Phe Asp Leu Asp Met Leu Gly Asp Gly Asp	
	175 180 185	
	TCC CCG GGG CCG GGA TTT ACC CCC CAC GAC TCC GCC CCC TAC GGC GCT	625
	Ser Pro Gly Pro Gly Phe Thr Pro His Asp Ser Ala Pro Tyr Gly Ala	
	190 195 200	

CTG GAT ATG GCC GAC TTC GAG TTT GAG CAG ATG TTT ACC GAT GCC CTT 673
Leu Asp Met Ala Asp Phe Glu Phe Glu Gln Met Phe Thr Asp Ala Leu
205 210 215

GGA ATT GAC GAG TAC GGT GGG TAGATCT 701
Gly Ile Asp Glu Tyr Gly Gly
220 225

10

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 226 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Lys Leu Leu Ser Ser Ile Glu Gln Ala Cys Asp Ile Cys Arg Leu
1 5 10 15
Lys Lys Leu Lys Cys Ser Lys Glu Lys Pro Lys Cys Ala Lys Cys Leu
20 25 30
Lys Asn Asn Trp Glu Cys Arg Tyr Ser Pro Lys Thr Lys Arg Ser Pro
35 40 45
Leu Thr Arg Ala His Leu Thr Glu Val Glu Ser Arg Leu Glu Arg Leu
50 55 60
Glu Gln Leu Phe Leu Leu Ile Phe Pro Arg Glu Asp Leu Asp Met Ile
65 70 75 80
Leu Lys Met Asp Ser Leu Gln Asp Ile Lys Ala Leu Leu Thr Gly Leu
85 90 95
Phe Val Gln Asp Asn Val Asn Lys Asp Ala Val Thr Asp Arg Leu Ala
100 105 110
Ser Val Glu Thr Asp Met Pro Leu Thr Leu Arg Gln His Arg Ile Ser
115 120 125
Ala Thr Ser Ser Ser Glu Glu Ser Ser Asn Lys Gly Gln Arg Gln Leu
130 135 140
Thr Val Ser Thr Ala Pro Pro Thr Asp Val Ser Leu Gly Asp Glu Leu
145 150 155 160
His Leu Asp Gly Glu Asp Val Ala Met Ala His Ala Asp Ala Leu Asp
165 170 175
Asp Phe Asp Leu Asp Met Leu Gly Asp Gly Asp Ser Pro Gly Pro Gly
180 185 190
Phe Thr Pro His Asp Ser Ala Pro Tyr Gly Ala Leu Asp Met Ala Asp
195 200 205
Phe Glu Phe Glu Gln Met Phe Thr Asp Ala Leu Gly Ile Asp Glu Tyr
210 215 220

60

Gly Gly
225

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GGCAACAATG AAGCTACTGT CTTCTATC

28

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

GGCAGATCTA CCCACCGTAC TCGTCAATTC C

31

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 109 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

GGCAAGCTTG GATCCAACAA TGAAGCTCCT GTCCTCCATC GAGCAGGCCT GCGACATCTG

60

CCGCCTCAAG AAGCTCAAGT GCTCCAAGGA GAAGCCGAAG TCGCCAAG

109

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 108 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

TCCTCTCGAG GGAAGATCAG GAGGAAGAGC TGCTCCAGGC GCTCCAGGCG GGACTCCACT 60
TCGGTGAGGT GGGCGCGGGT CAGCGGGGAG CGCTTGGTTT TGGGAGAG 108

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 81 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

TTCCCTCGAG AGGACCTCGA CATGATCCTG AAAATGGACT CCCTCCAGGA CATCAAAGCC 60
CTGCTCACCG GCCTCTTCGT C 81

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 89 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GGGTGAGGGG CATGTCGGTC TCCACGGAGG CCAGGCGGTC GGTGACGGCG TCTTTGTTCA 60
CGTTGTCCTG GACGAAGAGG CCGGTGAGC 89

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 80 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

GGAGACCGAC ATGCCCTCA CCCTGCGCCA GCACCGCATC AGCGCGACCT CCTCCTCGGA 60

GGAGAGCAGC AACAAAGGGCC

80

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 83 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

10	AGTGGAGCTC GTCCCCCAGG CTGACGTCGG TCGGGGGGGC CGTCGAGACG GTCAACTGGC	60
	GCTGGCCCTT GTTGCTGCTC TCC	83

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 56 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

20	ATATCCTGTC AAACACTGGA TCCGAGCTCC AATTCATAGT TTAAACTGAA GGCGGG	56
----	---	----

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

	GGCAGATCTT CGCAAGACCC TTCCTCTATA TAAGG	35
--	--	----

30

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

CACACAAACG GTGATACGTA

20

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 59 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

GAACTCTAGA AGCTACTCCA CGTCCATAAG GGACACATCA CAATCCCACT ATCCTTCGC

59

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 91 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

TAGGTTTACC CGCCAATATA TCCTGTCAAA CACTGGATCT TCGCAAGACC CTCCTCTAT

60

ATAAGGAAGT TCATTTTCATT TGGAGAGGAC A

91

CLAIMS

1. A nucleic acid sequence, expressible in a plant cell, encoding at least an effective portion of a GAL4 DNA-binding domain, the sequence having a %
5 A/T base content substantially reduced relative to the wild-type yeast sequence, characterised in that the % A/T base content is less than 45%.
2. A nucleic acid sequence is claimed in Claim 1 characterised in that the % A/T base content is less than 40%.
- 10 3. A sequence according to Claim 1 or Claim 2, further comprising a structural and/or regulatory sequence in operable linkage to the sequence encoding the effective portion of the GAL4 DNA-binding domain.
- 15 4. A sequence according to any one of Claims 1 to 3 wherein the nucleic acid encodes a peptide or polypeptide, having transcriptional activation activity, in operable linkage with the sequence encoding the effective portion of the GAL4 DNA-binding domain.
- 20 5. A sequence according to any one of Claims 1 to 4; wherein the sequence encodes a peptide or polypeptide, having transcriptional activation activity in a plant cell, which is a modified portion of GAL4, VPA16, GCN4 or a designed sequence which does not naturally occur.
- 25 6. A nucleic acid construct comprising a sequence in accordance with any one of claims 1 to 5.
7. A construct according to Claim 6, capable of inserting a sequence according to any one of Claims 1 to 5 into the genome of a plant host cell.
- 30 8. A construct according to Claim 6 or 7, further comprising a reporter gene in operable linkage with a GAL4-responsive upstream activation sequence (UAS).

AMENDED SHEET

- AMENDED SHEET

18. A method of expressing a gene of interest in a known pattern in a plant or part thereof, the method comprising: introducing the gene or interest into the plant or part thereof, said gene of interest having a GAL4- responsive upstream activation sequence (UAS), characterised in that said plant or part thereof
5 comprises a reporter gene expressed in a known pattern under the influence of a transcriptional activator comprising an effective portion of a GAL4 DNA-binding domain encoded by a sequence according to any one of Claims 1 to 5, such that binding of the transcriptional activator to the UAS causes transcriptional activation of the gene of interest.
- 10 19. A method according to Claim 18, wherein the gene of interest is introduced into one or more plants or parts thereof from a library according to any one of Claims 15, 16 or 17.
- 15 20. A method for co-ordinating the expression of a plurality of genes of interest in a plant or part thereof, each of the genes of interest being functionally associated with a GAL4-responsive UAS, characterised in that the plant or part thereof comprises a sequence according to any one of Claims 1 to 5 and is capable of
20 expressing a transcriptional activator comprising an effective portion of a GAL4 DNA-binding domain, such that binding of the transcriptional activator to the UAS causes simultaneous transcriptional activation of all of the genes of interest.
- 25 21. A method according to Claim 20, wherein at least some of the genes of interest are arranged polycistronically and associated with a single GAL4-responsive UAS.
22. A method according to Claim 20, wherein genes of interest are functionally associated with respective GAL4-responsive UASs.

AMENDED SHEET

WO 97/30164

PCT/GB97/00406

1/6

A/T	content
32/11	Hind III BamH I
60%	atg aag cta ctg tct tct atc gaa caa gca tgc gat att tgc
40%	<u>aaactt</u> <u>gaatcc</u> aaca atg aag ctg tcc tcc atc gag cag gcc tgc gac atc tgc
	M K L L S S I E Q A C D I C
92/31	
55%	62/21 cga ctt aaa aag ctc aag tgc tcc aaa gaa aaa ccg aag tgc gcc aag tgt ctg aag aac
45%	cgc ctc aag aag ctc aag tgc tcc aag gag aag ccg aag tgc gcc aag tgt ctg aag aac
	R L K K L K C S K E K P K C A K C L K N
152/51	
47%	122/41 aac tgg gag tgt cgc tac tct ccc aaa acc aaa agg tct ccg ctg act agg gca cat ctg
35%	aac tgg gag tgt cgc tac tct ccc aaa acc aaa agg cgc tcc ccg ctg acc cgc gcc cgc ctc
	N W E C R Y S P K T K R S P L T R A H L
212/71	Xho I
60%	182/61 aca gaa gtg gaa tca agg cta gaa aga ctg gaa cag cta ttt cta ctg att ttt cct cga
37%	acc gaa gtg gag tcc cgc ctg gag cgc ctg gag cag ctc ttc ctc ctg atc ttc cct cga
	T E V E S R L E R L E Q L F L L I F P R
272/91	
70%	242/81 gaa gac ctt gac atg att ttg aaa atg gat tct tta cag gat ata aaa gca ttg tta aca
42%	gag gac ctc gac atg atc ctg aaa atg gac tcc ctc ctc cag gac atc aaa gcc ctg ctc acc
	E D L D M I L K M D S L Q D I K A L L T
332/111	
63%	302/101 gga tta ttt gta caa gat aat gtg aat aaa gat gcc gtc aca gat aga ttg gct tca gtg
35%	ggc ctc ttc gtc cag gac aac gtg aac aaa gac gcc gtc acc gac cgc ctg gcc tcc gtg
	G L F V Q D N V N K D A V T D R L A S V

Fig. 1 Sheet 1

2/6

362/121 392/131 58%
 gag act gat atg cct cta aca ttg aga cag cat aga ata agt gcg aca tca tca tgc gaa
 gag acc gag atg ccc ctg acc ctg cgc cag cac cgc atc agc gcg acc tcc tcc tgc gag 30%
 E T D M P L T L R Q H R I S A T S S E
 422/141 452/151 47%
 gag agt agt aac aaa ggt caa aga cag ttg act gta tgc acg gcc ccc ccg acc gat gtc
 gag agc agc aac aag ggc cag cgc cag ttg acc gtc tgc acg gcc ccc ccg acc gag gtc 30%
 E S S N K G Q R Q L T V S T A P P T D V
 482/161 512/171 30%
 agc ctg ggg gac gag ctg cac tta gac ggc gag gac gtg gcg atg gcg cat gcc gac gcg
 agc ctg ggg gac gag ctg cac tta gac ggc gag gac gtg gcg atg gcg cat gcc gac gcg
 S L G D E L H L D G E D V A M A H A D A
 542/181 572/191 40%
 cta gac gat ttc gat ctg gac atg ttg ggg gac ggg gat tcc ccg ggg ccg gga ttt acc
 cta gac gat ttc gat ctg gac atg ttg ggg gac ggg gat tcc ccg ggg ccg gga ttt acc
 L D D F D L D M L G D G D S P G P G F T
 602/201 632/211 37%
 ccc cac gac tcc gcc ccc tac ggc gct ctg gat atg gcc gac ttc gag ttt gag cag atg
 ccc cac gac tcc gcc ccc tac ggc gct ctg gat atg gcc gac ttc gag ttt gag cag atg
 P H D S A P Y G A L D M A D F E F E Q M
 662/221 692/231 Bgl II
 ttt acc gat gcc ctt gga att gac gag tac ggt ggg tag
 ttt acc gat gcc ctt gga att gac gag tac ggt ggg tagatct
 F T D A L G I D E Y G G *

Fig. 1 Sheet 2

3/6




Fig. 2

WO 97/30164

PCT/GB97/00406

4/6

Modified Ti plasmid right border

TAGGTTTACCGCCCAATATATCCTGTCAAACACT GGATCTTCGCAAGACCCTT
|-----25 bp repeat-----|

mRNA start
|

CCTCTATATAAGGAAGTTCATTTTCAATTTGGAGAGGACA..GAL4 gene..
TATA Box

Fig. 3

WO 97/30164

PCT/GB97/00406

5/6

x
UAS-nGUS




Fig. 4a

X-gluc stained
brightfield

Fig. 4c

6/6



Fig. 4b



Fig. 4d